



PHY 2048 and 2049 formula sheet

$$v^2 = v_0^2 + 2a(x - x_0)$$

$$\sum F = ma$$

$$v=r\omega$$

$$a_c = \frac{v^2}{r}$$

$$p=mv$$

$$I=F\Delta t=\Delta p$$

$$I = \sum mr^2$$

$$\tau = I\alpha = r \cdot F$$

$$f = \frac{1}{2\pi} \sqrt{\frac{k}{m}}$$

$$f = \frac{1}{2\pi} \sqrt{\frac{g}{l}}$$

$$pAV=\text{constant}$$

$$p+\frac{1}{2}\rho v^2+\rho gy=\text{constant}$$

$$y=A\sin(kx-\omega t)$$

$$\omega = \frac{2\pi}{T} = 2\pi f$$

$$\beta = 10\log\left(\frac{l}{l_0}\right)$$

$$\frac{\Delta L}{L} = \alpha \Delta T \quad \frac{\Delta V}{V} = \beta \Delta T$$

Isothermal	Constant-volume	Isobaric	Adiabatic
$T=\text{constant}; Q=W$	$V=\text{constant}; Q=\Delta W$	$p=\text{constant}; Q=\Delta U+W$	$Q=0; \Delta U=-W$
$W=nRT\ln\left(\frac{V_2}{V_1}\right)$	$W=0$	$W=p(V_2-V_1)$	$W=\frac{p_1V_1-p_2V_2}{\gamma-1}$
$pV=\text{constant}$	$Q=nC_v\Delta T$	$Q=nC_p\Delta T; C_p=C_v+R$	$pV^\gamma = \text{constant}; TV^{\gamma-1} = \text{constant}$
$C_p = \frac{5}{2}R$	$C_v = \frac{3}{2}R$	$pV=nRT$	$dS = \frac{dQ}{T}$

This instructional aid was prepared by the Tallahassee Community College Learning Commons.



PHY 2048 and 2049 formula sheet

$$v^2 = v_0^2 + 2a(x - x_0)$$

$$\sum F = ma$$

$$v=r\omega$$

$$a_c = \frac{v^2}{r}$$

$$p=mv$$

$$I=F\Delta t=\Delta p$$

$$I = \sum mr^2$$

$$\tau = I\alpha = r \cdot F$$

$$f = \frac{1}{2\pi} \sqrt{\frac{k}{m}}$$

$$f = \frac{1}{2\pi} \sqrt{\frac{g}{l}}$$

$$pAV=\text{constant}$$

$$p+\frac{1}{2}\rho v^2+\rho gy=\text{constant}$$

$$y=A\sin(kx-\omega t)$$

$$\omega = \frac{2\pi}{T} = 2\pi f$$

$$\beta = 10\log\left(\frac{l}{l_0}\right)$$

$$\frac{\Delta L}{L} = \alpha \Delta T \quad \frac{\Delta V}{V} = \beta \Delta T$$

Isothermal	Constant-volume	Isobaric	Adiabatic
$T=\text{constant}; Q=W$	$V=\text{constant}; Q=\Delta W$	$p=\text{constant}; Q=\Delta U+W$	$Q=0; \Delta U=-W$
$W=nRT\ln\left(\frac{V_2}{V_1}\right)$	$W=0$	$W=p(V_2-V_1)$	$W=\frac{p_1V_1-p_2V_2}{\gamma-1}$
$pV=\text{constant}$	$Q=nC_v\Delta T$	$Q=nC_p\Delta T; C_p=C_v+R$	$pV^\gamma = \text{constant}; TV^{\gamma-1} = \text{constant}$
$C_p = \frac{5}{2}R$	$C_v = \frac{3}{2}R$	$pV=nRT$	$dS = \frac{dQ}{T}$

This instructional aid was prepared by the Tallahassee Community College Learning Commons.



PHY 2048 and 2049 formula sheet

$$\Delta E = Q + W$$

$$\vec{F} = \frac{kq_1 q_2}{r} \hat{r}$$

$$\vec{F} = q\vec{E}$$

$$\vec{E} = \frac{\vec{F}}{q_{test}} = \frac{kq}{r^2} \hat{r}$$

$$V(r) = \frac{kq}{r}$$

$$\oint \vec{E} \cdot d\vec{A} = \frac{q_{enclosed}}{\epsilon_0}$$

$$\frac{1}{c} = \frac{1}{c_1} + \frac{1}{c_2}$$

$$C = C_1 + C_2$$

$$V=IR$$

$$P=IV$$

$$P=I^2R$$

$$R=R_1+R_2$$

$$\vec{F} = q\vec{V} \times \vec{B}$$

$$\vec{F} = I\vec{L} \times \vec{B}$$

$$\oint \vec{B} \cdot d\vec{r} = \mu_0 I_{enclosed} \quad B = \mu_0 n I$$

$$B = \frac{\mu_0 NI}{2\pi r}$$

$$\varepsilon = -\frac{d\varphi_B}{dt} = BlV$$

$$\omega_0 = \frac{1}{\sqrt{LC}}$$

$$Z = \sqrt{R^2 + (X_L - X_C)^2}$$

$$X_C = \frac{1}{\omega C}$$

$$X_L = \omega L$$

$$n_1 \sin(\theta_1) = n_2 \sin(\theta_2)$$

$$\frac{1}{s} + \frac{1}{s'} = \frac{1}{f}$$

$$C = \frac{Q}{V}$$

$$V = V_c e^{-\frac{t}{RC}}$$

$$V_c = \epsilon(1 - e^{-\frac{t}{RC}}) \quad \phi = \vec{E} \cdot \vec{A} \quad \phi = \oint \vec{E} dA \cos \theta$$

This instructional aid was prepared by the Tallahassee Community College Learning Commons.



PHY 2048 and 2049 formula sheet

$$\Delta E = Q + W$$

$$\vec{F} = \frac{kq_1 q_2}{r} \hat{r}$$

$$\vec{F} = q\vec{E}$$

$$\vec{E} = \frac{\vec{F}}{q_{test}} = \frac{kq}{r^2} \hat{r}$$

$$V(r) = \frac{kq}{r}$$

$$\oint \vec{E} \cdot d\vec{A} = \frac{q_{enclosed}}{\epsilon_0}$$

$$\frac{1}{c} = \frac{1}{c_1} + \frac{1}{c_2}$$

$$C = C_1 + C_2$$

$$V=IR$$

$$P=IV$$

$$P=I^2R$$

$$R=R_1+R_2$$

$$\vec{F} = q\vec{V} \times \vec{B}$$

$$\vec{F} = I\vec{L} \times \vec{B}$$

$$\oint \vec{B} \cdot d\vec{r} = \mu_0 I_{enclosed} \quad B = \mu_0 n I$$

$$B = \frac{\mu_0 NI}{2\pi r}$$

$$\varepsilon = -\frac{d\varphi_B}{dt} = BlV$$

$$\omega_0 = \frac{1}{\sqrt{LC}}$$

$$Z = \sqrt{R^2 + (X_L - X_C)^2}$$

$$X_C = \frac{1}{\omega C}$$

$$X_L = \omega L$$

$$n_1 \sin(\theta_1) = n_2 \sin(\theta_2)$$

$$\frac{1}{s} + \frac{1}{s'} = \frac{1}{f}$$

$$C = \frac{Q}{V}$$

$$V = V_c e^{-\frac{t}{RC}}$$

$$V_c = \epsilon(1 - e^{-\frac{t}{RC}}) \quad \phi = \vec{E} \cdot \vec{A} \quad \phi = \oint \vec{E} dA \cos \theta$$

This instructional aid was prepared by the Tallahassee Community College Learning Commons.